

AN ODD THEOREM

BY

B. CURTIS EAVES

TECHNICAL REPORT NO. 69-10
SEPTEMBER 1969

Operations Research House
Stanford University
Stanford, California

Research and reproduction of this report was partially supported by Office of Naval Research, Contracts ONR-N-00014-67-A-0112-0011 and ONR-N-00014-67-A-0112-0016; U.S. Atomic Energy Commission, Contract AT [04-3] -326 PA #18; National Science Foundation Grant GP-6431; U.S. Army Research Office, Contract DAHCO4-67-C0028; and National Institute of Health, Grant GM 14789-02.

Reproduction in whole or in part is permitted for any purpose of the United States Government. This document has been approved for public release and sale; its distribution is unlimited.

AD699893

AN ODD THEOREM

BY

B. CURTIS EAVES

TECHNICAL REPORT NO. 69-10
SEPTEMBER 1969

Operations Research House
• Stanford University
• Stanford, California

Research and reproduction of this report was partially supported by Office of Naval Research, Contracts ONR-N-00014-67-A-0112-0011 and ONR-N-00014-67-A-0112-0016; U.S. Atomic Energy Commission, Contract AT[04-3]-326 PA #18; National Science Foundation Grant GP-6431; U.S. Army Research Office, Contract DAHCO4-67-C0028; and National Institute of Health, Grant GM 14789-02.

Reproduction in whole or in part is permitted for any purpose of the United States Government. This document has been approved for public release and sale; its distribution is unlimited.

Abstract

Let C be a bounded convex polyhedral set and let $f:C \rightarrow C$ be continuous and piecewise linear. Using notions from complementary pivot theory, it is shown that if each fixed point of f lies interior to some piece of linearity, then f has an odd number of fixed points. In addition, an algorithm is given for computing a fixed point of f .

1. Introduction

Using the main ideas of complementary pivot theory (see [1] - [8]), we prove the following theorem.

Theorem: Let C be a bounded convex polyhedral set and let $f:C \rightarrow C$ be continuous and piecewise linear. If each fixed point of f lies interior to some piece of linearity, then f has an odd number of fixed points.

An algorithm for computing (finitely quick) a fixed point of f (whether or not the interior condition is met) is a by-product of the proof of the theorem.

The essential difference between our attitude and that of [1], [4], [8], and hence Sperner's Lemma is that we label vertices of a triangulation with vectors instead of integers. For a simplex to be "completely labeled," there must be a convex combination of the vector labels which generate zero.

2. Graph Principle

Our proof will rest on a simple graph principle; the same principle used in [1] - [8]. By a graph (N,A) , we mean a finite set N together with a symmetric anti-reflexive relation A on N . If aAb (hence bAa and $a \neq b$), we say a and b are adjacent. We call an element a of N odd or even if it is adjacent to an odd or even number of elements of N , respectively. Recall that a graph has an even number of odd elements. In the next section, we construct a particular graph and use this device to prove our theorem. In this graph each element will be adjacent to exactly one or two elements; in this case, the odd elements have a natural pairing.

3. The Theorem and Proof

Let C be a finite dimensional bounded convex polyhedral set. We can assume that C lies in n -dimensional Euclidean space and that it has an interior. Let T be a triangulation of C (i.e., T is a complex and $|T| = C$, see [9]), and let $f: C \rightarrow C$ be a continuous function which is linear (i.e., affine) on each simplex of T . Let (C, T, f) denote such a triple. If each fixed point of f is interior to an n -simplex of T , then we say that (C, T, f) is nondegenerate. To prove our result, it is sufficient to prove the following theorem.

Theorem: If (C, T, f) is nondegenerate, then f has an odd number of fixed points.

Given (C, T, f) we notice that f is completely determined by its action on the vertices of T . Indeed, if $r \in S \in T$, then $f(r) = \sum_s f(s)x_s$ where $r = \sum_s sx_s$, $\sum_s x_s = 1$, and $x_s \geq 0$ (where s ranges over the vertices of S).

A simplex S of T contains a fixed point if and only if the system

$$\sum_s (f(s) - s)x_s = 0$$

$$\sum_s x_s = 1$$

has a nonnegative solution in the x_s (where s ranges over the vertices of S). In this case $\sum_s sx_s$ is the fixed point. If (C, T, f) is nondegenerate, then it follows that a simplex will contain at most one fixed point; if S contains a fixed point, then S is an n -simplex and the solution x_s of the system above is unique and positive.

Assume (C, T, f) is nondegenerate. Let C' be an n -simplex which contains C in its interior. We shall extend both T and f to C' to generate (C', T', f') . Let v_0, \dots, v_n be the vertices of C' .

Extend T to a triangulation T' of C' without introducing new vertices; that is, a vertex of T' is either a vertex of T or a vertex of C' . Hence each $(n-1)$ -face of C' is an element of T' .

Temporarily let r be any point of C . Define f' on C' by setting $f'(t) = f(t)$ for vertices of T and $f'(v_i) = r$ for vertices of C' and then by extending f' linearly on the simplexes of T' . Now we further specify r . Select $r \in C$ such that for any $(n-1)$ -simplex S of T' the system


$$\begin{aligned} (f'(v_0) - v_0)x_S + \sum_s (f'(s) - s)x_s &= 0 \\ x_S + \sum_s x_s &= 1 \end{aligned}$$

either has a unique positive solution in x_S and the x_s or else has no non-negative solution (where s ranges over the vertices of S). Such r 's are very available; in fact, almost every element of C will suffice.

We can now define a particular graph. Let (C', T', f') be generated as just described. Let N_1 be the set of simplexes of T' which contain fixed points; these simplexes will be n -simplexes of T . Let N_2 be the set of $(n-1)$ -simplexes S in T' for which the system

$$\begin{aligned} (f'(v_0) - v_0)x_S + \sum_s (f'(s) - s)x_s &= 0 \\ x_S + \sum_s x_s &= 1 \end{aligned}$$

has a nonnegative solution in x_S and the x_s (where s ranges over the vertices of S); these solutions will be positive and unique. Let $N = N_1 \cup N_2$. We define two distinct simplexes of N to be adjacent if they lie in a common simplex of T' .



Let S_0 be the $(n-1)$ -simplex with vertices $\{v_1, \dots, v_n\}$. One can now establish that $S_0 \in N_2$, that each element of $N_1 \cup \{S_0\}$ is adjacent to exactly one element of N , and that each element of $N_2 \setminus \{S_0\}$ is adjacent to exactly two elements of N . From the graph principle, we see that N_1 contains an odd number of elements; this establishes the theorem.

The figure illustrates the structure for a 2-dimensional C . The arrow at a vertex t denotes the direction of $f'(t)-t$ (further specification is unnecessary), the heavy lines denote the boundary of C , and the small circles denote the simplexes which are in N .

4. The Algorithm

The preceding development gives a procedure for calculating finitely quick a fixed point of (C, T, f) . After constructing (C', T', f') , one begins at S_0 and proceeds to an adjacent simplex, etc. This step from simplex to simplex is essentially a "pivot" as known in linear programming. One eventually terminates with a simplex containing a fixed point, and hence, one has the fixed point.

The next section shows that if (C, T, f) is degenerate, the algorithm may still be applied to find a fixed point ((C, T, f) is altered slightly to make it nondegenerate; however, from solely computational considerations, there are far more efficient methods of dealing with degeneracy).

The section on Brouwer's Theorem demonstrates that if $g: C \rightarrow C$ is a continuous function and if $\epsilon > 0$, then the algorithm can be used to compute a point $t \in C$ such that $|g(t) - t| \leq \epsilon$. Scarf's procedure [8] has this capability.

Kuhn [4] describes an extremely efficient data handling procedure which can be adapted to our algorithm.

5. Perturbation and Stability

Here we show that nondegeneracy is stable and that nondegeneracy can be achieved via a perturbation.

Suppose that (C, T, f) is nondegenerate. Then there is an $\epsilon > 0$ such that (S, T, g) is nondegenerate and such that a simplex of T will contain a fixed point of f if and only if it contains a fixed point of g , if $|f - g| \leq \epsilon$.

Consider (C, T, f) and (C, T, g) . Suppose that $g(C) = r$ and that r is interior to an n -simplex of T . Then there is an $\epsilon_0 > 0$ such that for $0 < \epsilon \leq \epsilon_0$, $(C, T, (1-\epsilon)f + \epsilon g)$ is nondegenerate. Further, if a simplex of T contains a fixed point of $(1-\epsilon)f + \epsilon g$, then it contains a fixed point of f for $0 < \epsilon \leq \epsilon_0$.

6. Brouwer's Theorem and Extensions

From Sections 4 and 6 we see that for any (C, T, f) there is a fixed point. We can now prove Brouwer's fixed point theorem.

Let $g: C \rightarrow C$ be a continuous function. Choose (C, T_n, f_n) such that $|f_n - g| \leq \frac{1}{n}$ for $n=1, 2, \dots$. Let s_n be a fixed point of f_n . We have $|g(s_n) - s_n| \leq \frac{1}{n}$. If s is a cluster point of the s_n sequence, then clearly s is a fixed point of g .

For the general case $g: C \rightarrow C$ where g is continuous and C is compact and convex, our theorem has implications regarding the parity of the number of fixed points. These results will be reported on in another paper.

References

1. Cohen, D. I. A., "On the Sperner Lemma," J. Combinatorial Theory 2, 585-7 (1967).
2. Cottle, R. W. and G. B. Dantzig, "Complementary Pivot Theory of Math. Programming," Linear Algebra and Its Applications, 1, 103-125 (1968).
3. Eaves, B. Curtis, "Linear Complementarity Problem in Math. Programming," Doctoral Dissertation, Stanford University, 1969.
4. Kuhn, Harold W., "Simplicial Approximation of Fixed Points," Proceedings of the National Academy of Sciences of the U.S.A., 16, 1237-42 (1968).
5. Lemke, C. E., "Bimatrix Equilibrium Points and Mathematical Programming," Management Science, 11, 681-689 (1965).
6. Lemke, C. E., and J. T. Howson, Jr., "Equilibrium Points of Bimatrix Games," SIAM J. Appl. Math., 12, 413-423 (1964).
7. Murty, K. G., "On the Number of Solutions to the Complementary Quadratic Programming Problem," Doctoral Dissertation, Engineering Science, University of California, Berkeley, 1968.
8. Scarf, H., "The Approximation of Fixed Points of a Continuous Mapping," SIAM J. Appl. Math., 15, 1328-43 (1967).
9. Pontryagin, L. S., "Foundations of Combinatorial Topology," Graylock Press, Rochester, New York, 1952.

Unclassified
Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Department of Operations Research Stanford University Stanford, Calif		2a. REPORT SECURITY CLASSIFICATION
		2b. GROUP
3. REPORT TITLE AN ODD THEOREM		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Technical Report		
5. AUTHOR(S) (Last name, first name, initial) EAVES, B. Curtis		
6. REPORT DATE September 1969	7a. TOTAL NO. OF PAGES 6	7b. NO. OF REFS 9
8a. CONTRACT OR GRANT NO. N00014-67-A-0112-0011	9a. ORIGINATOR'S REPORT NUMBER(S) 69-10	
b. PROJECT NO. NR-047-064	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
c.		
d.		
10. AVAILABILITY/LIMITATION NOTICES This document has been approved for public release and sale: its distribution is unlimited		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Operations Research Program (code 434) Office of Naval Research Washington, D. C.	
13. ABSTRACT Let C be a bounded convex polyhedral set and let $f:C \rightarrow C$ be continuous and piecewise linear. Using notions from complementary pivot theory, it is shown that if each fixed point of f lies interior to some piece of linearity, then f has an odd number of fixed points. In addition, an algorithm is given for computing a fixed point of f .		

DB FORM 1 JAN 64 1273

Unclassified
Security Classification